



# Imidacloprid 2F

## *Turf & Ornamental Insecticide*

### FOLIAR AND SYSTEMIC INSECT CONTROL

For use on turfgrass (including sod farms), landscape ornamentals, fruit and nut trees, and interior plantscapes. *For use by professional personnel licensed or registered by the state to apply turf maintenance, nursery/greenhouse, and/or landscaping/ornamental products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the structural pest control regulatory agency of your state prior to use of this product.*

#### ACTIVE INGREDIENT:

	% BY WEIGHT
Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine .....	21.4%
INERT INGREDIENTS: .....	78.6%
TOTAL: .....	100.0%

Contains 2 pounds of imidacloprid per gallon. Shake well before using.

EPA Reg. No. 53883-229-73220

EPA Est. No. 53883-TX-002

## KEEP OUT OF REACH OF CHILDREN

# CAUTION

PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

Manufactured for: FarmSaver.com, LLC • 4515 Falls of Neuse Road, Suite 300 • Raleigh, NC 27609

## Net Contents: 1 Gallon

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of soap and water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eyelids open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Prosar at 1-800-308-5391 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN</b>	
No specific antidote is available. Treat the patient symptomatically.	

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets away from treated area until dry.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

#### **WPS Uses:**

Applicators and other handlers (mixers and loaders) who handle this product for uses covered by the Worker Protection Standard (40 CFR part 170) – such as sod must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **Non-WPS Uses:**

- Applicators and other handlers must wear:
- Shirt and pants
- Gloves
- Shoes plus socks

**ENGINEERING CONTROLS STATEMENTS:** When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(4)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### **USER SAFETY RECOMMENDATIONS**

#### **Users should:**

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Remove and wash contaminated clothing before reuse.

### **ENVIRONMENTAL HAZARDS**

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Apply this product only as specified on this label. Extreme care must be taken to avoid runoff. Apply only to soil or other fill substrate that will accept the solution at the specified rate. Do not treat soil that is water-saturated or frozen or in any conditions where run-off or movement from the treatment area (site) is likely to occur.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

(continued)

### **AGRICULTURAL USE REQUIREMENTS (continued)**

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

**Exception:** If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton. if you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.
- Shoes plus socks

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

### **APPLICATION ON TURFGRASS**

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE may be used to control insect pests on turfgrass in home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields, and sod farms.

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE controls soil inhabiting pests such as Northern & Southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotrogus majalis*; Green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; Billbugs *Sphenophorus* spp.; Annual bluegrass weevil, *Hyperodes* spp.; Black turfgrass ataenius,

Ataenius spretulus and Aphodius spp. European Crane Fly Tipula paludosa, and mole crickets, scapteriscus spp. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE can also be used for suppression of cutworms and chinch bugs.

For optimum control, make applications preceding or during the egg laying period of the target pest. The active ingredient in QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE has enough residual activity so that applications can be made preceding the egg laying activity. Application timing can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Most favorable control will be achieved when applications are made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Do not make applications when turfgrass areas are waterlogged or the soil is saturated with water. Sufficient distribution of the active ingredient cannot be achieved under these conditions. The treated turf area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Applications cannot exceed a total of 1.6 pints (0.4 lb of active ingredient) per acre per year.

#### **APPLICATION EQUIPMENT FOR USE ON TURFGRASS**

Apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for the application of turfgrass insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.

Do not apply through any irrigation system.

## RECOMMENDED APPLICATIONS

### TURF GRASSES

PEST	RATE	APPLICATION INSTRUCTIONS
Larvae of: Annual bluegrass weevil, Asiatic garden beetle, Billbug, Black turfgrass ataenius, Cutworms (suppression), European chafer, European Crane Fly, Green June beetle, Japanese beetle, Northern Masked chafer, Oriental beetle, <i>Phyllophaga</i> spp., Southern masked chafer	1 .25 to 1.6 pt/A or 0.46 to 0.6 fl. oz. (14 to 17 mL) per 1,000 sq. ft.	For best control of grubs, billbugs, annual bluegrass weevil, and European Crane Fly, apply prior to egg hatch of the target pest. Read <b>APPLICATION EQUIPMENT</b> section of this label.
Chinchbugs (suppression) Mole crickets	1.6 pt /A or 0.6 fl. oz. (17 mL) per 1,000 sq. ft.	For suppression of chinchbugs, apply before hatching of the first instar nymphs.  To control mole crickets apply before or during the peak egg hatch period. Use a curative insecticide in addition to QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE when adults or large nymphs are present and actively tunneling. Follow label instructions for other insecticides when tank-mixing.
Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialists for more specific information regarding timing of application.		
<b>NOTE:</b> For best control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. Do not apply more than 1.6 pt (0.4 lb of active ingredient) per acre per year. Avoid mowing turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.		

### APPLICATION TO ORNAMENTALS

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE is for use on ornamentals in commercial and residential landscapes and interior plantscapes. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE is a systemic product and will be taken up into the plant system from root uptake. To assure optimum effectiveness, the product must be placed where the grow-

ing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution may enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests.

When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is taken up throughout the plant. In some cases, this translocation delay could take 60 days or longer. For this reason, applications should be made prior to anticipated pest infestation to achieve optimum levels of control.

For outdoor ornamentals, broadcast applications cannot exceed a total of 1.6 pints (0.4 lb of active ingredient) per acre per year.

### **Ant Management Programs**

Use QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE to control aphids, scale insects, mealybugs and other sucking pests on ornamentals to limit the honeydew available as a food source for ant populations. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE applications can be then be supplemented with residual sprays, bait placements or other ant control tactics to further reduce the pest population.

**Note:** *Not for use in commercial greenhouses, nurseries, or on grasses grown for seed, or on commercial fruit and nut trees.*

### **APPLICATION EQUIPMENT FOR FOLIAR APPLICATIONS**

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE mixes readily with water and may be used in many types of application equipment. Mix product with the required amount of water and apply as desired dependent upon the selected use pattern.

When making foliar applications on hard to wet foliage such as holly, pine, or ivy, the addition of a spreader/ sticker is recommended. If concentrate or mist type spray equipment is used, an equivalent amount of product should be used on the area sprayed, as would be used in a dilute application.

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE has been found to be compatible with commonly used fungicides, miticides, liquid fertilizers, and other commonly used insecticides. Check physical compatibility using the correct proportion of products in a small jar test if local experience is unavailable.

Do not apply through any irrigation system.



**RECOMMENDED APPLICATIONS****FOR USE ONLY IN AND AROUND INDUSTRIAL AND COMMERCIAL BUILDINGS AND RESIDENTIAL AREAS.**

<b>CROP</b>	<b>PEST</b>	<b>RATE</b>	<b>APPLICATION INSTRUCTIONS</b>
Trees Shrubs Evergreens Flowers Foliage plants Groundcovers Interior plantscapes	Adelgids Aphids Japanese beetles Lace bugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy winged sharpshooter) Mealybugs Psyllids Sawfly Larvae Thrips (suppression) Whiteflies	1.5 fl. oz. (45 mL) per 100 gal of water	<b>Foliar Applications:</b> Begin applications before the onset of high pest populations and reapply as needed.
	White grub larvae (such as Japanese beetle larvae, Chafer, Phyllophaga spp. Asiatic garden beetle, Oriental beetle)	0.46 to 0.6 fl. oz. (14 to 17 mL) per 1,000 sq. ft.	<b>Broadcast Applications:</b> Use enough water to mix the product and thoroughly apply to the treatment area. Do not use less than 2 gallons of water per 1,000 sq ft. For best results, irrigate after application to incorporate QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE into the upper soil layer.  For additional use directions, refer to the <b>FLOWERS and GROUND COVERS</b> section of this label.

# SOIL INJECTION\* AND SOIL DRENCH APPLICATIONS IN AND AROUND INDUSTRIAL AND COMMERCIAL BUILDINGS AND RESIDENTIAL AREAS, AND STATE, NATIONAL, AND PRIVATE WOODED AND FORESTED AREAS

\*No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.

PEST	CROP/RATE	APPLICATION INSTRUCTIONS	REMARKS
<p>Adelgids</p> <p>Aphids</p> <p>Armored scales (suppression)</p> <p>Black vine weevil larvae</p> <p>Emerald ash borer</p> <p>Eucalyptus longhorned borer</p> <p>Flatheaded borer (including bronze birch and alder borer)</p> <p>Japanese beetles</p> <p>Lace bugs</p> <p>Leaf beetles (including elm and viburnum leaf beetles)</p> <p>Leafhoppers (including glassy-winged sharpshooter)</p> <p>(continued)</p>	<p><b>TREES</b></p> <p>0.1 to 0.2 fl. oz. (3 to 6 mL) per inch of trunk diameter (D.B.H.)</p>	<p><b>SOIL INJECTION:</b></p> <p><b>Grid System:</b> Space holes in a grid pattern on 2.5 foot centers, extending to the drip line of the tree.</p> <p><b>Circle System:</b> Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line.</p> <p><b>Basal System:</b> Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.</p> <p><b>Soil Drench:</b> Apply uniformly as a drench around the base of the tree in not less than 10 gallons of water per 1,000 square feet. Direct application to the root area. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p>	<p>Use enough water to mix the product and inject an equal amount of solution in each hole. Use low pressure and sufficient solution for distribution of the liquid into the treatment area. For best control, keep the treated area moist for 7 to 10 days.</p> <p>Do not use less than 4 holes per tree.</p> <p><b>For Control of Specified Borers:</b> Trees with existing insect damage and stress may not recover after treatment with QUALI-PRO IMIDA-CLOPRID 2F TURF &amp; ORNAMENTAL INSECTICIDE.</p>

PEST (continued)	CROP/RATE	APPLICATION INSTRUCTIONS	REMARKS
Leafminers Mealybugs Pine tip moth larvae Psyllids Royal palm bugs Sawfly larvae Soft scales Thrips (suppression) White grub larvae Whiteflies	<b>SHRUBS</b>  0.1 to 0.2 fl. oz. (3 to 6 mL) per foot of shrub height	<b>Soil Injection:</b> Apply at the recommended dosage to each plant.  <b>Soil Drench:</b> Apply uniformly as a drench around the base of the tree in not less than 10 gallons of water per 1000 square feet. Direct application to the root area. Remove plastic or any other barrier that will stop solution from reaching the root zone.	Use enough water to mix the product and inject an equal amount of solution in each hole. Use low pressure and sufficient solution for distribution of the liquid into the treatment area. For best control, keep the treated area moist for 7 to 10 days.  Do not use less than 4 holes per shrub.
	<b>FLOWERS AND GROUND COVERS</b>  0.46 to 0.6 fl. oz. (14 to 17 mL) per 1,000 sq ft	Apply as a broadcast treatment before or after planting, or apply after plants are established. Mix into soil. For best control on established plants, irrigate thoroughly after application.	

# FOLIAR APPLICATIONS FOR USE ONLY IN AND AROUND RESIDENTIAL AREAS

CROP	PEST	RATES	APPLICATION INSTRUCTIONS
<b>POME FRUITS</b> Apple Crabapple Loquat Mayhaw Pear Pear (oriental) Quince	Aphids (except Woolly apple aphid) Leafhoppers (including glassy-winged sharpshooter) Leafminer Mealybugs* San Jose scale*	1.5 fl. oz. (45 mL) per 100 gal or 6.0 fl. oz./A <sup>1</sup>	<p>Apply specified dosage as foliar spray as needed after petal-fall is complete.</p> <p>For control of rosy apple aphid, apply prior to leafrolling caused by the pest.</p> <p>For first generation leafminer control, make first application as soon as petal-fall is complete. Greatest leafminer control will result from the earliest possible application.</p> <p>For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. QUALI-PRO IMIDACLOPRID 2F TURF &amp; ORNAMENTAL INSECTICIDE will not control late stage larvae.</p> <p>For San Jose Scale, time applications to the crawler stage. Treat each generation.</p> <p>For late season (preharvest) control of leafhopper species, apply QUALI-PRO IMIDACLOPRID 2F TURF &amp; ORNAMENTAL INSECTICIDE while most leafhoppers are in the nymphal stage.</p> <p>For optimal control of mealybug, insure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.</p> <p>Do not apply more than 6.0 fluid ounces per acre in a single application. Do not make more than 5 applications.</p> <p>Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.</p> <p>* Not for use in California for control on pears.</p>

(continued)

**FOLIAR APPLICATIONS FOR USE ONLY IN AND AROUND RESIDENTIAL AREAS (continued)**

CROP	PEST	RATES	APPLICATION INSTRUCTIONS
Pecan*	Yellow pecan aphid Black margined aphid Pecan leaf phylloxera Pecan spittlebug Pecan stem phylloxera	1.5 fl. oz. (45 mL) per 100 gal or 6.0 fl. oz./A <sup>1</sup>	<p>Make foliar applications as pests begin to build before populations become extreme. Two applications at a 10 to 14 day interval may be required to achieve control. Scout and retreat if needed.</p> <p>Thorough uniform coverage of foliage is necessary for optimal control. Addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's recommended use rate may improve coverage.</p> <p>Do not apply more than a total of 18.0 fluid ounces of QUALI-PRO IMIDACLOPRID 2F TURF &amp; ORNAMENTAL INSECTICIDE per acre per year. Do not make more than 3 applications.</p> <p>Allow 10 or more days between applications.</p> <p>* Use on pecans not permitted in California unless directed by specific supplemental labeling.</p>
<sup>1</sup> The amount of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.			

**FOLIAR APPLICATION FOR USE ONLY IN AND AROUND INDUSTRIAL AND COMMERCIAL BUILDINGS AND RESIDENTIAL AREAS.**

<b>CROP</b>	<b>PEST</b>	<b>RATES</b>	<b>APPLICATION INSTRUCTIONS</b>
Grapes	Leafhoppers (including glassy-winged sharpshooter) Mealybugs	1.5 fl. oz. (45 mL) per 100 gal or 3.0 fl. oz/A (90 mL/A)	Apply specified dosage as a foliar spray using 200 gallons of water per acre. Do not apply more than a total of 6.0 ounces of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE per acre per year.  Allow at least 14 days between applications. Applications may be applied up to and including day of harvest.

**RESTRICTIONS**

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Avoid runoff or puddling of irrigation water following application. Keep children and pets off treated area until dry.
- Avoid application of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE to areas which are water logged or saturated, which will not allow penetration into the root zone of the plant.
- Do not apply more than 1.6 pt (0.4 lb of active ingredient) per acre per year.

Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store water soluble packets in original container and out of reach of children, preferably in a locked storage area.

*(continued)*

### STORAGE AND DISPOSAL (continued)

Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticide below. In spill or leak incidents, keep unauthorized people away.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site (in the treatment area) or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### IMPORTANT: READ BEFORE USE

#### LIMITATION OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of FarmSaver.com, LLC. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, FarmSaver.com, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of FarmSaver.com, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, FarmSaver.com, LLC disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all

losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at FarmSaver.com, LLC's election, the replacement of product.

Quali-Pro is a registered trademark of FarmSaver.com, LLC.

EPA app 07-20-07



Glue.